## Original Article

## Peer Reviewed





International Journal of Experimental Research and Review (IJERR)

© Copyright by International Academic Publishing House (IAPH)

ISSN: 2455-4855 (Online)

www.iaph.in



Screening for Chronic Obstructive Pulmonary Disease in Patients with Coronary Artery Disease at a **Tertiary Care Hospital** 

Check for updates

Malasree Jujjavarapu, Meenakshi Narasimhan\*, Sridhar Rathinam and Shahana Balasubramanian

Department of Respiratory Medicine, Chettinad Hospital and Research Institute, Chettinad Academy of Research and Education, Kelambakkam-603103, Tamil Nadu, India

#### E-mail/Orcid Id:

MJ, @ malasreej96@gmail.com, https://orcid.org/0009-0007-4465-8815; MN, paddy\_2020@yahoo.com, https://orcid.org/0000-0002-8743-149X; SR, @ srihema.1964@gmail.com, 10 https://orcid.org/0000-0001-7661-2351; SB, 12 drshahana93@gmail.com, 10 https://orcid.org/0009-0001-8753-4413

#### **Article History**:

Received: 26th May, 2024 Accepted: 17th July, 2024 Published: 30th July, 2024

#### **Keywords:**

COPD, CAD, Spirometry, **PRISM** 

#### How to cite this Article:

Malasree Jujjavarapu, Meenakshi Narasimhan, Sridhar Rathinam and Shahana Balasubramanian (2024). Screening for Chronic Obstructive Pulmonary Disease in Patients with Coronary Artery Disease at a Tertiary Care Hospital. International Journal of Experimental Research and Review, 41(spl.), 120-129.

https://doi.org/10.52756/ijerr.2024.v41spl.010

Abstract: Chronic Obstructive Pulmonary Disease is a common, preventable and treatable disease that is characterized by persistent respiratory symptoms and airflow limitation that is due to airway and/or alveolar abnormalities usually caused by significant exposure to noxious gases or particles and influenced by host factors, including abnormal lung development. COPD was the 4th most common cause of death globally, and it was anticipated to rise to the 3rd position by 2020 or sooner. Over 80% of these patients go undiagnosed. COPD and CAD frequently co-exist and interact due to various reasons like similar risk factors and pathogenesis. Although there are global studies on the prevalence of COPD in CAD patients, data from India are scarce. This study aims to find the prevalence of COPD among CAD patients and find the correlation between the occurrence of COPD and cardiac status. This study was conducted as a prospective observational cross-sectional study at Chettinad Hospital and Research Institute in India. 293 patients with angiographically proven CAD underwent pulmonary function evaluation by spirometry to assess the prevalence of COPD. Recent 2D-ECHO was also correlated. The study found that 18.77% of 293 Coronary Artery Disease (CAD) patients had COPD and 3.07% had PRISM( preserved ratio impaired spirometry, a new term coined), making it one of the few studies to report the prevalence of PRISM among CAD patients. Outside of India, studies have found that COPD prevalence ranges from 7% to 33.6% among ischemic heart disease patients. Impairment of spirometry positively correlated with the severity of CAD. Our study found a considerable frequency of missed COPD diagnoses. Hence, we recommend that when CAD is diagnosed, the initial evaluation should include a complete history of respiratory symptoms, clinical examination, and spirometry to assess pulmonary function, particularly in those who have risk factors such as smoking and the use of biomass fuel so as to identify COPD at an early stage and treat it. Identifying PRISM can help prevent its progression to COPD by making the required lifestyle modifications. A multidisciplinary approach is advised for all those diagnosed to have CAD.

## Introduction

COPD will continue to become more common worldwide (Murray et al., 1997; Sarkar et al., 2021; Tamondong-Lachica et al., 2023; Dange et al., 2023). COPD was the 4th most common cause of death globally, and it was anticipated to rise to the 3rd position by 2020 or sooner(Murray et al., 1997; Lopez et al., 2006). COPD is frequently missed in India due to a lack of spirometry

testing and symptoms that appear gradually. The prevalence of COPD varies depending on the study technique and country, but among persons 40 years and older, it ranges from 4% to 20% (Halbert et al., 2006; Buist et al., 2007; Adeloye et al., 2022). The prevalence of Chronic Obstructive Pulmonary Disease in India is 7.4% (Onishi et al., 2014).

Cardiovascular disorders, diabetes, COPD and cancer are among India's most frequent chronic illnesses (Madhu et al., 2022, 2023; Sur et al., 2013; Biswas et al., 2023). As people embrace sedentary behaviours and bad diets, the prevalence of CVDs, diabetes, COPD and cancer is projected to rise (Roy et al., 2023; Sarkar et al., 2023; Paritala et al., 2024). Furthermore, improved diagnostic methods and increased awareness of these disorders lead to higher identification rates (Yach et al., 2004). More than 80% of these individuals go untreated, increasing their risk of cardiovascular complications (Buist et al., 2007; Halbert et al., 2003). Cigarette smoking is a major risk factor for COPD however, not every smoker develops it. Cigarette smoke has also been linked to the development of cardiovascular disease (CVD), which is another leading cause of death worldwide (Yach et al., 2004).

It has been argued that the risk of CVD increases in smokers who develop COPD and COPD could be a risk factor for developing CVD (Sin et al., 2005; Sin et al., 2008). However, the precise prevalence of COPD in CVD patients remains unknown, and there has been little research on COPD screening in CAD patients (Rodriguez-Roisin et al., 2008).

A study conducted in Spain found airflow limitation in 19% of CVD patients (Soriano et al., 2010), whereas airflow limitation was observed in 10%-25% of patients visiting a university hospital's cardiovascular department in Japan (Wada et al., 2010; Yamasaki et al., 2010). However, the risk of comorbid COPD and CVD is poorly understood in India.

COPD and CAD frequently co-exist, adding to the risk of morbidity and mortality. The presence and progression of CAD is one of the causes of COPD exacerbations, which can result in major cardiovascular events, including cardiovascular death.

The coexistence of chronic obstructive pulmonary

disease and coronary artery disease is complex and not fully understood. Both these disorders have similar risk factors, namely- smoking, age, diet, sedentary lifestyle and air pollution as well as similar disease pathways like inflammation, hypoxia and oxidative stress (Madhu et al., 2023; Prasad et al., 2023). These factors contribute to the development and progression of both conditions.

inflammatory Smoking induces responses in susceptible individuals, which promote inflammation (Maiti and Samanta, 2018; Dey and Guha, 2020). This inflammation is associated with COPD and formation, worsening and rupture atherosclerotic plaques. These processes can cause coronary artery disease, heart failure, and mortality. Additionally, the hypoxia experienced by COPD patients also contributes to the development of CAD. As a result, screening for COPD in CAD can lead to early intervention and optimal COPD care, resulting in fewer complications and deaths. Therefore, a cross-sectional study was undertaken using spirometry to identify COPD in Coronary Artery Disease patients. This study intends to determine the prevalence of COPD among CAD patients and the relationship between COPD and the patients' underlying cardiac condition.

#### **Materials and Methods**

This study was a prospective observational cross-sectional study at Chettinad Hospital and Research Institute in India. All Coronary artery disease (CAD) patients diagnosed by Coronary angiogram attending Cardiology OPD were enrolled into the study after fulfilling inclusion criteria and obtaining written consent. Their demographic details and clinical details were collected through a structured proforma. After obtaining cardiac fitness, all participants were subjected to detailed clinical examination, imaging by Chest X-ray, ECG, 2D-Echo, and spirometry with bronchodilator reversibility.




Figure 1. Outline of the research methodology

Study diagnosis was based on guidelines developed by the Global Initiative for Chronic Obstructive Lung Disease (GOLD, 2023).

## Sample Size And Sampling Technique

A convenience sampling technique was used for the study. 293 patients who visited the Cardiology OPD at CHRI, Kelambakkam and met the inclusion criteria were enrolled on the study.

#### **Inclusion Criteria**

- 1) Patients with stable coronary artery disease diagnosed by coronary angiogram.
  - 2) Age: 35-80 years.
  - 3) Gender: All genders.

#### **Exclusion Criteria**

- 1) Patients who do not consent to the study.
- 2) Patients with contraindications for spirometry as per ATS/ERS guidelines
- 3)Patients who are previously diagnosed to have COPD.

Table 1. Sex distribution.

Gender	Frequency	Percentage
Male	178	60.7%
Female	115	39.24%

## **Ethical permission**

Ethical committee approval was obtained at our institute to carry on this study (IHEC- I/1103/22)

#### **Statistical analysis**

SPSS software, version 25.0, was used to perform statistical analyses.

#### Result

## **Profile Of Patients**

We enrolled 293 patients diagnosed with coronary artery disease (CAD) through coronary angiography at our center who met the inclusion criteria. The mean age

observed among patients was 55.65 years ( $\pm 8.37$ ). Maximum patients were in the age group 47–57 years (46.07%) (Figure 2). 178 patients (60.7%) were males and 115 (39.3%) were females (Table 1). 138 patients (47.09%) in the study group had normal BMI, 0.6% were underweight, 39.24% were overweight and 12.96% were obese.

21.84 % were smokers, 39.93% were former smokers and 15.6% had exposure to biomass fuel. All the smokers were males.

## **Symptoms**

Amongst the subjects, dyspnoea was the most common symptom (48.1%), followed by cough (13%), breathlessness and cough (3%) and 44.3% of the subjects were asymptomatic (Figure 3).

#### **Chest Xray P/A View:**

(81.5%) Chest X-ray was normal, whereas hyperinflation was seen in 11%, cardiomegaly-3.75% and increased broncho vascular markings-3.75% (Figure 4).

# Cardiac Status Based On Coronary Angiogram And 2D-ECHO

56.99% of patients had minimal CAD, 10.58% had moderate CAD and 32.4% had severe CAD (figure 5). On echocardiography, 80.54% of patients had a normal left ventricular ejection fraction (LVEF > 55%), 12.28% had LVEF between 50 and 55% and 7.1% had LVEF between 41 and 49%, none had LVEF below 40% (Figure 6). Pulmonary Hypertension was present in 7.84% of the cases, whereas 92.15% had no pulmonary hypertension.

#### **COPD Among CAD Patients**

COPD was diagnosed on spirometry when postbronchodilator FEV1/FVC < 0.7.

PRISM was diagnosed when postbronchodilator FEV1/FVC>0.7, but other spirometry parameters were abnormal.




Figure 2. Age distribution.

••




Figure 3. Symptomology.




Figure 4. Chest x-ray findings.




Figure 5. Severity of CAD as per coronary angiogram.




Figure 6. Ejection fraction.

After spirometry, out of 293 patients in the study group, 55 patients (18.77%) were diagnosed to have COPD and 9 patients (3.07%) were diagnosed as PRISM (Preserved ratio impaired spirometry)(Figure 6). In a survey of 5487 female adults over 45 years old, the authors found a 7.1% prevalence of PRISm, similar to findings (Wijnant et al., 2020). epidemiological research has revealed the prevalence of PRISm is 5.1% in the general population and 12.3 percent among smokers (Wan et al., 2021; Nonato et al., 2015). 229 patients (78.1%) had a normal spirometry. Amongst COPD 32(58.2%) were male and 23(41.8%) were females. Amongst the PRISM candidates- 4(44.4%) were males and 5(55.5%) were females. Among the COPD cases, 10.90% had mild COPD, 89.09% had moderate COPD, and none had severe or severe COPD (Figure 7). Amongst the COPD cases, 30 patients were smokers and 25 patients had biomass fuel exposure.

# Association Between Severity Of CAD And Spirometry

We assessed the correlation between the interpretation of spirometry and CAD Severity. According to the chisquare value of 32.47, the P-value was found to be 0.0001. This was statistically correlated and determined to be highly significant. Impairment of spirometry positively correlated with the severity of CAD.

#### **Discussion**

The patients had an average age of 55.65 years (±8.37), similar to findings reported in prior research conducted in India and internationally (Rosenberg et al., 2015). Most patients (46.07%) were aged 47 to 57 years. The majority (60.7%) of patients were male, which is a similar finding reported by two prior studies(Mahendra et al., 2018; Agarwal et al., 2019; Balbirsingh et al., 2023; Wijnant et al., 2020; Ferreira et al., 2023). It is fair, given that CAD and tobacco smoking are more common among men. Dyspnea was reported by 48% of patients. Cough symptoms were reported in roughly 4% of the cases. In 3% of cases, cough and dyspnea occurred simultaneously. 44% of the cases were asymptomatic. According to (Alisherovna et al., 2023; Mota et al., 2018; Mannino et al., 2005), cough and dyspnea were present in all cases. This is similar to our study.




Figure 7. Spirometry interpretation.

In our study, 82% of the patients demonstrated normal chest X-rays. Atypical x-ray features, such as increased broncho vascular markings, occur in around 3.75% of cases. Cardiomegaly was found in 3.75%. 10.93% had overinflated lung fields. Daher et al. (2020) did a recent study that highlighted the findings of chest X-ray observations. According to the study, around 78% of all cases had normal chest X-ray results. However, it was shown that 5% of cases had cardiomegaly (enlarged heart) and 17% had pleural effusion (Daher et al., 2020).

Tobacco usage is linked to both cardiovascular and obstructive airway diseases. In our study, approximately 21.84% of patients were smokers, while 39.93% were former smokers. This is smaller than that reported by Indian-71% (Mahendra et al.,2018) and international studies-72.6% (Khassawneh et al., 2018).

echocardiographic findings of tricuspid regurgitation, right ventricular enlargement, right atrial enlargement, and pulmonary arterial hypertension. However, none of our research participants experienced tricuspid regurgitation, right ventricular enlargement, or right atrial enlargement. As COPD worsens, these disorders become more widespread, and the risk of cardiac dysfunction rises.

Out of 293 patients with stable CAD, 55 (18.77%) had COPD on spirometry and 9 (3.07%) had PRISM. Outside of India, studies have found that COPD prevalence ranges from 7% to 33.6% among ischemic heart disease patients(Behar et al., 1992; Wada et al., 2010).) In another study with a large number of patients (995), the prevalence of airflow limitation (AL) was reported to be 27%, although pulmonary function testing (PFT) was




Figure 8. Severity of COPD.

The coronary angiography revealed that the majority of the patients had mild to severe coronary disease. On angiography, the majority of patients (56.99%) had mild CAD, whereas 10.58% had moderate CAD and 32.4% had severe CAD. Out of 293 instances, 80.54% had a normal ejection fraction, while 12.29% showed diastolic dysfunction. 7.17% experienced both systolic and diastolic dysfunction. Nobody had isolated systolic dysfunction. A large percentage of people exhibited diastolic dysfunction, while a lesser number had mixed dysfunction. In the current investigation, about 7.84% of cases had pulmonary hypertension.

Similar to our study (Ansari et al., 2024), the study sought to determine how varying levels of COPD severity influenced echocardiographic results. They concluded that there is a high link between COPD severity and done using a handheld spirometer and a FEV1/FEV6 < 0.73 was defined as AL (Onishi et al., 2014; Verma et al., 2021). In our study, FEV1/FVC <0.7 post-bronchodilator was considered as airflow limitation. Previous research has found comparable results for underdiagnosis and undertreatment of COPD in patients with ischemic heart disease. (Mahendra et al., 2018; Khassawneh et al., 2018). Ours is one of the few studies to report the incidence of PRISM amongst CAD cases.

## Our study had certain limitations like:

# Patients with severe COPD and cardiac complications who could not perform Spirometry could not be included in the study.

# The actual prevalence of COPD in CAD patients could have been more accurate had we included the known cases of COPD.

# The study had a small sample size and was conducted at a single location, so the findings cannot be generalised to the entire community.

These limitations can be overcome in future studies by enrolling even the known cases of COPD who have been diagnosed as having CAD to know the real prevalence of COPD amongst CAD patients. A larger sample size may be considered for future investigations.

#### **Conclusion**

COPD treatment enhances the quality of life and reduces future cardiovascular risk and mortality. Early detection of the presence of obstructive airway illness as soon as the patient's cardiac condition stabilises and timely treatment will result in significant reductions in morbidity and mortality over time. Screening and early detection of COPD in CAD can lead to earlier intervention and optimal COPD care, resulting in fewer complications and deaths. Effective management of COPD and CAD requires a multidisciplinary approach. Collaboration among respiratory and cardiovascular specialists is critical for providing complete care. This missed diagnosis of COPD and PRISM is quite common. Identifying PRISM is especially crucial because these patients can develop COPD in the future.

Lifestyle changes made at the time of PRISM identification may help prevent the development of COPD. We recommend that at the time of CAD diagnosis, the initial evaluation includes a complete history, clinical examination, and spirometry to assess lung function, particularly in patients with risk factors such as smoking and biomass fuel use. Patients diagnosed with COPD should be counselled for smoking cessation and the start of appropriate treatment, as reduction of airway limitation improves cardiovascular health.

Therefore, we conclude that once a patient is diagnosed with CAD by a cardiologist, once stabilised, the patient may be referred to the respiratory medicine physician to analyse the pulmonary function status further so as to identify the undiagnosed COPD and PRISM. A multidisciplinary approach is advised.

#### Acknowledgement

I would like to thank the Department of Respiratory Medicine and Cardiology at CHRI, Kelambakkam, for giving me the opportunity to undertake this study. I sincerely like to thank my faculty and colleagues for their constant encouragement and guidance. My love to my family and friends for being my strength.

## **Conflict of Interest**

The authors declare no conflict of interest.

#### References

Adeloye, D., Song, P., Zhu, Y., Campbell, H., Sheikh, A., Rudan, I., & NIHR RESPIRE Global Respiratory Health Unit (2022). Global, regional, and national prevalence of, and risk factors for, chronic obstructive pulmonary disease (COPD) in 2019: a systematic review and modelling analysis. *The Lancet. Respiratory Medicine*, *10*(5), 447–458. https://doi.org/10.1016/S2213-2600(21)00511-7

Agrawal, A., Singh, S. K., & Prabhudesai, P. P. (2019). The prevalence of obstructive airway disease by spirometric indices in non-smoker patients with catheter diagnosed coronary artery disease. *Indian J. Respir. Care*, 8, 46-50.

https://doi.org/10.4103/ijrc.ijrc\_31\_18

Alisherovna, K.M., Akramovna, I.K., Bakhtiyorovich, U. J., Nizamitdinovich, K.S., Jasurovna, J. S., Kairatovna, R. A., & Abdukholikovna, E. S. (2023). Exacerbations of chronic obstructive pulmonary disease and coronary atherosclerosis. *Journal of New Century Innovations*, *39*(1), 176-178.

Ansari, R. N., Prajapati, V. B., Chavda, S. H., & Kapadia, R. C. Cardiac Assessment of Chronic Obstructive Pulmonary Disease Patients by ECHO and Its Association with Different Grades of Severity of Chronic Obstructive Pulmonary Disease.

Balbirsingh, V., Mohammed, A. S., Turner, A. M., & Newnham, M. (2022). Cardiovascular disease in chronic obstructive pulmonary disease: a narrative review. *Thorax*, thoraxjnl-2021-218333. *Advance online publication*. https://doi.org/10.1136/thoraxjnl-2021-218333

Behar, S., Panosh, A., Reicher-Reiss, H., Zion, M., Schlesinger, Z., & Goldbourt, U. (1992). Prevalence and prognosis of chronic obstructive pulmonary disease among 5,839 consecutive patients with acute myocardial infarction. SPRINT Study Group. *The American Journal of Medicine*, 93(6), 637–641. https://doi.org/10.1016/0002-9343(92)90196-i

Biswas, T., Behera, B. K., & Madhu, N.R. (2023). Technology in the Management of Type 1 and Type 2 Diabetes Mellitus: Recent Status and Future Prospects. 26 pages, Springer Nature Singapore Pte Ltd., Advances in Diabetes Research and Management. pp. 111–136. Online ISBN-978-981-19-0027-3. https://doi.org/10.1007/978-981-19-0027-3\_6

Buist, A. S., McBurnie, M. A., Vollmer, W. M.,Gillespie, S., Burney, P., Mannino, D. M., Menezes,A. M., Sullivan, S. D., Lee, T. A., Weiss, K. B.,Jensen, R. L., Marks, G. B., Gulsvik, A.,

Nizankowska-Mogilnicka, E., & BOLD Collaborative Research Group (2007). International variation in the prevalence of COPD (the BOLD Study): a population-based prevalence study. *Lancet (London, England)*, *370*(9589), 741–750.

https://doi.org/10.1016/S0140-6736(07)61377-4

Daher, A., & Dreher, M. (2020). The bidirectional relationship between chronic obstructive pulmonary and coronary artery disease. disease bidirektionale Beziehung zwischen chronisch obstruktiver Lungenerkrankung und koronarer Herzkrankheit. Herz, 45(2), 110-117. https://doi.org/10.1007/s00059-020-04893-4

Dange, H., Patil, R., & Upasani, C. (2023). A Clinical Study of Allergy and COPD Correlation With Spirometric and FENO Evaluation. *Int. J. Exp. Res. Rev.*, *34*(Special Vol.), 138-145.

https://doi.org/10.52756/ijerr.2023.v34spl.013

- Dey, S., & Guha, P. (2020). A brief review on cigarette induced cellular. *Int. J. Exp. Res. Rev.*, 23, 18-26. https://doi.org/10.52756/ijerr.2020.v23.002
- Ferreira, E. J. P., Cardoso, L. V. S. C., de Matos, C. J. O., Mota, I. L., Lira, J. M. C., Lopes, M. E. G., Santos, G. V., Dória Almeida, M. L., Aguiar-Oliveira, M. H., Sousa, A. C. S., de Melo, E. V., & Oliveira, J. L. M. (2023). Cardiovascular Prognosis of Subclinical Chronic Obstructive Pulmonary Disease in Patients with Suspected or Confirmed Coronary Artery Disease. *International Journal of Chronic Obstructive Pulmonary Disease*, 18, 1899–1908. https://doi.org/10.2147/COPD.S410416
- Halbert, R. J., Isonaka, S., George, D., & Iqbal, A. (2003). Interpreting COPD prevalence estimates: what is the true burden of disease? *Chest*, *123*(5), 1684–1692. https://doi.org/10.1378/chest.123.5.1684
- Halbert, R. J., Natoli, J. L., Gano, A., Badamgarav, E., Buist, A. S., & Mannino, D. M. (2006). Global burden of COPD: systematic review and meta-analysis. *The European respiratory journal*, 28(3), 523–532.

https://doi.org/10.1183/09031936.06.00124605.

- Khassawneh, B. Y., Samrah, S. M., Jarrah, M. I., Ibdah, R. K., Ibnian, A. M., Al-Mistarehi, A. W., Zghayer, A. A., Abuqudairi, S. I., & Khader, Y. S. (2018). Prevalence of undiagnosed COPD in male patients with coronary artery disease: a cross-sectional study in Jordan. *International Journal of Chronic Obstructive Pulmonary Disease*, 13, 2759–2766. https://doi.org/10.2147/COPD.S172679
- Lopez, A. D., Shibuya, K., Rao, C., Mathers, C. D., Hansell, A. L., Held, L. S., Schmid, V., & Buist, S.

(2006). Chronic obstructive pulmonary disease: current burden and future projections. *The European Respiratory Journal*, 27(2), 397–412.

https://doi.org/10.1183/09031936.06.00025805

- Madhu, N.R., Sarkar, B., Biswas, P., Roychoudhury, S., Behera, B.K., & Acharya, C.K. (2023). Therapeutic potential of melatonin in glioblastoma: Current knowledge and future prospects. Biomarkers in Cancer Detection and Monitoring of Therapeutics, Volume-2. Elsevier Inc., pp. 371-386. ISBN 978-0-323-95114-2. https://doi.org/10.1016/B978-0-323-95114-2.00002-9
- Madhu, N.R., Sarkar, B., Roychoudhury, S., Behera, B.K. (2022). Melatonin Induced in Cancer as a Frame of Zebrafish Model. © Springer Nature Singapore Pte Ltd. 2022, S. Pathak et al. (eds.), Handbook of Animal Models and its Uses in Cancer Research, pp. 1-18. ISBN: 978-981-19-1282-5.

https://doi.org/10.1007/978-981-19-1282-5\_61-1

- Madhu, N.R., Sarkar, B., Slama, P., Jha, N.K., Ghorai, S.K., Jana, S.K., Govindasamy, K., Massanyi, P., Lukac, N., Kumar, D., Kalita, J.C., Kesari, K.K., & Roychoudhury, S. (2022). Effect of Environmental Stressors, Xenobiotics, and Oxidative Stress on Male Reproductive and Sexual Health. © The Author(s), under exclusive license to Springer Nature Switzerland AG 2022, S. Roychoudhury, K. K. Kesari (eds.), Oxidative Stress and Toxicity in Reproductive Biology and Medicine. *Advances in Experimental Medicine and Biology*, *1391*, 33-58. ISBN: 978-3-031-12966-7.
  - https://doi.org/10.1007/978-3-031-12966-7\_3.
- Mahendra, M., S, S. K., Desai, N., Bs, J., & Pa, M. (2018). Evaluation for airway obstruction in adult patients with stable ischemic heart disease. *Indian Heart Journal*, 70(2), 266–271.

https://doi.org/10.1016/j.ihj.2017.08.003

- Maiti, M., & Samanta, G. (2018). Relationship between physical activity and smoking behavior among college students. *Int. J. Exp. Res. Rev.*, *15*, 39-43. https://doi.org/10.52756/ijerr.2018.v15.006
- Mannino, D. M., Holguin, F., Pavlin, B. I., & Ferdinands, J. M. (2005). Risk factors for prevalence of and mortality related to restriction on spirometry: findings from the First National Health and Nutrition Examination Survey and follow-up. The international journal of tuberculosis and lung disease: the official journal of the International Union against Tuberculosis and Lung Disease, 9(6), 613–621.

- Mota, I. L., Sousa, A. C. S., Almeida, M. L. D., de Melo, E. V., Ferreira, E. J. P., Neto, J. B., Matos, C. J. O., Telino, C. J. C. L., Souto, M. J. S., & Oliveira, J. L. M. (2018). Coronary lesions in patients with COPD (Global Initiative for Obstructive Lung Disease stages I-III) and suspected or confirmed coronary arterial disease. *International Journal of Chronic Obstructive Pulmonary Disease*, 13, 1999–2006. https://doi.org/10.2147/COPD.S162713
- Murray, C. J., & Lopez, A. D. (1997). Alternative projections of mortality and disability by cause 1990-2020: Global Burden of Disease Study. *Lancet (London, England)*, 349(9064), 1498–1504. https://doi.org/10.1016/S0140-6736(96)07492-2.
- Nonato, N. L., Nascimento, O. A., Padilla, R. P., de Oca, M. M., Tálamo, C., Valdivia, G., Lisboa, C., López, M. V., Celli, B., Menezes, A. M., & Jardim, J. R. (2015). Occurrence of respiratory symptoms in persons with restrictive ventilatory impairment compared with persons with chronic obstructive pulmonary disease: The PLATINO study. *Chronic Respiratory Disease*, 12(3), 264–273. https://doi.org/10.1177/1479972315588004
- Onishi, K., Yoshimoto, D., Hagan, G. W., & Jones, P. W. (2014). Prevalence of airflow limitation in outpatients with cardiovascular diseases in Japan. *International journal of chronic Obstructive Pulmonary Disease*, *9*, 563–568. https://doi.org/10.2147/COPD.S59962.
- Paritala, A., Lakshmanan, M., Narasimhan, M., & Rathinam, S. (2024). Evaluation of Pulmonary Status of Post-Tuberculosis Patients with Spirometry and Chest X-Ray. *International Journal of Experimental Research and Review*, 40(Spl Volume), 227-234. https://doi.org/10.52756/ijerr.2024.v40spl.019
- Prasad, N., Bhattacharya, T., & Lal, B. (2023). Chemometric Techniques in the Assessment of Ambient Air Quality and Development of Air Quality Index of Coal Mining Complex: A Statistical Approach. *Int. J. Exp. Res. Rev.*, 36, 433-446. https://doi.org/10.52756/ijerr.2023.v36.018a
- Rodríguez-Roisin, R., & Soriano, J. B. (2008). Chronic obstructive pulmonary disease with lung cancer and/or cardiovascular disease. *Proceedings of the American Thoracic Society*, *5*(8), 842–847. https://doi.org/10.1513/pats.200807-075TH.
- Rosenberg, S. R., Kalhan, R., & Mannino, D. M. (2015). Epidemiology of Chronic Obstructive Pulmonary Disease: Prevalence, Morbidity, Mortality, and Risk Factors. *Seminars in Respiratory and Critical Care*

- *Medicine*, 36(4), 457–469. https://doi.org/10.1055/s-0035-1555607
- Roy, R., Chakraborty, A., Jana, K., Sarkar, B., Biswas, P., & Madhu, N.R. (2023). The Broader Aspects of Treating Diabetes with the Application of Nanobiotechnology. Springer Nature Singapore Pte Ltd., Advances in Diabetes Research and Management, pp. 137–162, Online ISBN-978-981-19-0027-3.
  - https://doi.org/10.1007/978-981-19-0027-3 7
- Sarkar, B., Biswas, P., Acharya, C.K., Ghorai, S.K., Nahar, N., Jana, S.K., Ghosh, S., Sarkar, D., Behera, B., & Madhu, N.R. (2021). Knowledge of Traditional Indian Medicinal Plants for the Management of COPD. *Chettinad Health City Medical Journal*. 10(4), 184 189. https://doi.org/10.36503/chcmj10(4)-05
- Sarkar, S., Sadhu, S., Roy, R., Tarafdar, S., Mukherjee, N., Sil, M., Goswami, A., & Madhu, N.R. (2023). *Contemporary Drifts in Diabetes Management. Int. J. App. Pharm.*, 15(2), 1-9.
  - https://doi.org/10.22159/ijap.2023v15i2.46792
- Sin, D. D., & Man, S. F. (2005). Chronic obstructive pulmonary disease as a risk factor for cardiovascular morbidity and mortality. *Proceedings of the American Thoracic Society*, 2(1), 8–11.
  - https://doi.org/10.1513/pats.200404-032MS
- Sin, D. D., & Man, S. F. (2008). Impact of cancers and cardiovascular diseases in chronic obstructive pulmonary disease. *Current Opinion in Pulmonary Medicine*, *14*(2), 115–121.
  - https://doi.org/10.1097/MCP.0b013e3282f45ffb
- Soriano, J. B., Rigo, F., Guerrero, D., Yañez, A., Forteza, J. F., Frontera, G., Togores, B., & Agustí, A. (2010). High prevalence of undiagnosed airflow limitation in patients with cardiovascular disease. *Chest*, *137*(2), 333–340. https://doi.org/10.1378/chest.09-1264
- Sur, T., Das, A., Bashar, S., Tarafdar, S., Sarkar, B., & Madhu, N.R. (2023). Biochemical Assay for Measuring Diabetes Mellitus. Springer Nature Singapore Pte Ltd., Advances in Diabetes Research and Management, pp. 1–20, Online ISBN-978-981-19-0027-3.
  - https://doi.org/10.1007/978-981-19-0027-3\_1
- Tamondong-Lachica, D. R., Skolnik, N., Hurst, J. R., Marchetti, N., Rabe, A. P. J., Montes de Oca, M., & Celli, B. R. (2023). GOLD 2023 Update: Implications for Clinical Practice. *International Journal of Chronic Obstructive Pulmonary Disease*, 18, 745–754.
  - https://doi.org/10.2147/COPD.S404690

- Verma, A., Gudi, N., Yadav, U. N., Roy, M. P., Mahmood, A., Nagaraja, R., & Nayak, P. (2021). Prevalence of COPD among population above 30 years in India: A systematic review and metaanalysis. *Journal of Global Health*, 11, 04038. https://doi.org/10.7189/jogh.11.04038
- Wada, H., Nakano, Y., Nagao, T., Osawa, M., Yamada, H., Sakaguchi, C., Matsumoto, T., Tsutamoto, T., Ito, M., & Horie, M. (2010). Detection and prevalence of chronic obstructive pulmonary disease in a cardiovascular clinic: evaluation using a hand held FEV<sub>1</sub>/FEV<sub>6</sub> meter and questionnaire. *Respirology (Carlton, Vic.)*, 15(8), 1252–1258.

https://doi.org/10.1111/j.1440-1843.2010.01854.

Wan, E. S., Balte, P., Schwartz, J. E., Bhatt, S. P., Cassano, P. A., Couper, D., Daviglus, M. L., Dransfield, M. T., Gharib, S. A., Jacobs, D. R., Jr, Kalhan, R., London, S. J., Navas-Acien, A., O'Connor, G. T., Sanders, J. L., Smith, B. M., White, W., Yende, S., & Oelsner, E. C. (2021). Association Between Preserved Ratio Impaired Spirometry and Clinical Outcomes in US Adults. *JAMA*, 326(22), 2287–2298.

https://doi.org/10.1001/jama.2021.20939

- Wijnant, S. R. A., De Roos, E., Kavousi, M., Stricker, B. H., Terzikhan, N., Lahousse, L., & Brusselle, G. G. (2020). Trajectory and mortality of preserved ratio impaired spirometry: the Rotterdam Study. *The European Respiratory Journal*, *55*(1), 1901217. https://doi.org/10.1183/13993003.01217-2019
- Yach, D., Hawkes, C., Gould, C. L., & Hofman, K. J. (2004). The global burden of chronic diseases: overcoming impediments to prevention and control. *JAMA*, 291(21), 2616–2622. https://doi.org/10.1001/jama.291.21.2616
- Yamasaki, A., Hashimoto, K., Hasegawa, Y., Okazaki, R., Yamamura, M., Harada, T., Ito, S., Ishikawa, S., Takami, H., Watanabe, M., Igishi, T., Kawasaki, Y., & Shimizu, E. (2010). COPD is frequent in conditions of comorbidity in patients treated with various diseases in a university hospital. *International Journal of Chronic Obstructive Pulmonary Disease*, 5, 351–355. https://doi.org/10.2147/COPD.S12669

#### How to cite this Article:

Malasree Jujjavarapu, Meenakshi Narasimhan, Sridhar Rathinam and Shahana Balasubramanian (2024). Screening for Chronic Obstructive Pulmonary Disease in Patients with Coronary Artery Disease at a Tertiary Care Hospital. *International Journal of Experimental Research and Review*, *41*(spl.), 120-129.

**DOI**: https://doi.org/10.52756/ijerr.2024.v41spl.010



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.